

Physics formula list class xii

Physics Formula List Class XII: The Complete Guide Physics formula list class XII is an essential resource for students preparing for board exams, competitive exams, and understanding core concepts in physics. Having a comprehensive list of formulas helps students to revise quickly, grasp the fundamentals, and solve problems efficiently. This article provides an extensive compilation of physics formulas categorized topic-wise, ensuring that students can find the information they need in one place, along with explanations and tips for mastering them.

Important Physics Formulas in Class XII

1. Kinematics Kinematics deals with the motion of objects without considering the forces causing the motion. The formulas are fundamental for solving problems related to velocity, acceleration, and displacement.

Displacement with constant acceleration: $s = ut + \frac{1}{2}at^2$ Final velocity: $v = u + at$ Velocity squared: $v^2 = u^2 + 2as$ Average velocity (for uniformly accelerated motion): $v_{avg} = \frac{u + v}{2}$

2. Laws of Motion Newton's laws form the foundation of classical mechanics. The formulas here are crucial for understanding forces and motion.

Force (Newton's Second Law): $F = ma$ Weight: $W = mg$ Impulse: $J = F \Delta t = \Delta p$ Momentum: $p = mv$

3. Work, Power, and Energy This section includes formulas related to energy conservation, work done by forces, and power.

Work done: $W = F s \cos \theta$ Kinetic energy: $KE = \frac{1}{2}mv^2$ Potential energy (due to gravity): $PE = mgh$ Power: $P = \frac{W}{t}$

4. Rotational Motion Understanding rotational dynamics is vital for topics involving spinning objects and angular quantities.

Angular displacement: θ (radians) Angular velocity: $\omega = \frac{\theta}{t}$ Angular acceleration: $\alpha = \frac{\omega - \omega_0}{t}$ Moment of inertia (for a point mass): $I = mr^2$ Torque: $\tau = I\alpha$

5. Gravitation Gravitational formulas are key to understanding planetary motion, satellite dynamics, and gravitational fields.

Universal law of gravitation: $F = G \frac{m_1 m_2}{r^2}$ Acceleration due to gravity: $g = G \frac{M}{r^2}$ Gravitational potential energy: $U = -G \frac{m_1 m_2}{r}$ Escape velocity: $v_e = \sqrt{\frac{2GM}{r}}$

6. Properties of Matter This section covers formulas related to elasticity, stress, strain, and surface tension.

Hooke's Law: $F = kx$ Stress: $\sigma = \frac{F}{A}$ Strain: $\epsilon = \frac{\Delta l}{l}$ Young's modulus: $Y = \frac{\sigma}{\epsilon}$ Surface tension (force per unit length): $T = \frac{F}{l}$

7. Thermodynamics Formulas here are crucial for understanding heat, work, and efficiency in thermodynamic processes.

First Law of Thermodynamics: $\Delta U = Q - W$ Specific heat capacities (constant volume and pressure): C_v, C_p Work done in expansion/compression: $W = P \Delta V$ Carnot efficiency: $\eta = 1 - \frac{T_C}{T_H}$

8. Electrostatics Electrostatic formulas help in understanding electric charges, fields, and potentials.

Electric force (Coulomb's Law): $F = k_e \frac{q_1 q_2}{r^2}$ Electric field: $E = \frac{F}{q}$ or for point charges: $E = k_e \frac{q}{r^2}$ Electric potential: $V = \frac{U}{q}$ Potential difference: $V = Ed$

9. Current Electricity Formulas related to electric current, resistance, and circuits are fundamental in electronics.

Ohm's Law: $V = IR$ Power dissipated in a resistor: $P = VI = I^2 R = \frac{V^2}{R}$ Series resistance: $R_{total} = R_1 + R_2 + \dots + R_n$ Parallel resistance: $\frac{1}{R_{total}} = \frac{1}{R_1} + \frac{1}{R_2} + \dots + \frac{1}{R_n}$

10. Magnetism and Electromagnetism This section covers formulas related to magnetic fields, forces,

and electromagnetic induction. Magnetic force on a moving charge: $F = q v B \sin \theta$ Magnetic field due to a long straight current-carrying wire: $B = \frac{\mu_0 I}{2 \pi r}$ Electromagnetic induction (Faraday's Law): $\mathcal{E} = - \frac{d \Phi_B}{dt}$ Self-induction (inductive reactance): $X_L = 2 \pi f L$

Physics Formula List Class XII: A Comprehensive Guide for Students and Enthusiasts

Understanding the fundamental principles of physics hinges greatly on memorizing and mastering key formulas that describe the natural world. For Class XII students, who are preparing for board exams and beyond, having a well-organized, detailed compilation of physics formulas is essential. This article aims to serve as a thorough reference, breaking down complex concepts into digestible, analytical explanations, and emphasizing the significance of each formula within its respective domain.

Introduction to Physics Formulas for Class XII

Physics, often regarded as the backbone of understanding the universe, is a subject rooted in mathematical formulations. These formulas encapsulate the laws governing motion, energy, electricity, magnetism, optics, and modern physics. Mastery over these formulas not only aids in solving numerical problems efficiently but also deepens conceptual understanding.

Class XII physics syllabus encompasses a broad spectrum of topics, each with its own set of fundamental formulas. This guide aims to systematically explore these formulas, their derivations, applications, and the underlying principles they represent.

Kinematics

Kinematics deals with the motion of objects without considering the causes of motion. It introduces equations that describe displacement, velocity, acceleration, and time.

- Equations of Motion** These equations are valid for uniformly accelerated motion in a straight line:
 - First Equation: $v = u + at$ Where: (v) = final velocity (u) = initial velocity (a) = acceleration (t) = time elapsed
 - Second Equation: $s = ut + \frac{1}{2} a t^2$ Where: (s) = displacement
 - Third Equation: $v^2 = u^2 + 2as$
 These equations are foundational in analyzing linear motion, allowing students to determine unknown quantities when others are known. They are derived from calculus-based principles but are often used in their algebraic forms.
- Newton's Laws of Motion** Newton's laws form the core of classical mechanics, describing the relationship between forces and motion.
 - First Law (Law of Inertia)** An object remains at rest or moves uniformly in a straight line unless acted upon by an external force. This law introduces the concept of inertia and sets the stage for understanding force.
 - Second Law** $F = ma$ Where: (F) = net force applied on an object (m) = mass of the object (a) = acceleration produced. This formula quantifies how forces influence motion. It helps in calculating the acceleration resulting from a known force or vice versa.
 - Third Law** For every action, there is an equal and opposite reaction.
- Work, Energy, and Power** These concepts describe how energy is transferred and transformed in physical systems.
 - Work Done** $W = F s \cos \theta$ Where: (W) = work done (F) = magnitude of force (s) = displacement (θ) = angle between force and displacement. The work-energy principle states that the work done on an object results in a change in its energy.
 - Kinetic Energy (KE)** $KE = \frac{1}{2} m v^2$ This formula quantifies the energy possessed by a moving body.
 - Potential Energy (PE)** $PE = mgh$ Where: (m) = mass (g) = acceleration due to gravity (h) = height above a reference point.
 - Power** $P = \frac{W}{t}$

$\frac{W}{t}$ Expresses the rate at which work is done or energy is transferred.

Gravitation The study of forces of attraction between masses is central to understanding celestial mechanics and planetary motion.

1. Universal Law of Gravitation $[F = G \frac{m_1 m_2}{r^2}]$ Where: (F) = gravitational force between two masses (G) = gravitational constant $(6.674 \times 10^{-11} \text{ Nm}^2/\text{kg}^2)$ (m_1, m_2) = masses (r) = distance between centers of the two masses This law explains gravity's inverse-square nature and underpins planetary orbits and satellite motion.

2. Acceleration Due to Gravity $[g = \frac{GM}{r^2}]$ Where: (M) = mass of the celestial body (r) = radius from the center of the body to the point of measurement

Properties of Matter and Fluids This section deals with the behavior of solids, liquids, and gases under various forces.

1. Pressure and Buoyancy Pressure at a Depth (h) : $[P = P_0 + \rho g h]$ Where: (P_0) = atmospheric pressure (ρ) = density of the fluid Archimedes' Principle: $(F_b) = \rho_{\text{fluid}} \times g \times V_{\text{displaced}}$ This explains why objects float or sink.

2. Bernoulli's Theorem $[P + \frac{1}{2} \rho v^2 + \rho g h = \text{constant}]$ Describes the behavior of flowing fluids, illustrating the relationship between pressure, velocity, and height.

Thermodynamics The study of heat, work, and energy transfer.

1. Specific Heat Capacity $[Q = mc \Delta T]$ Where: (Q) = heat energy supplied (m) = mass (c) = specific heat capacity (ΔT) = change in temperature

2. Gas Laws Boyle's Law: $[PV = \text{constant}]$ Charles's Law: $[\frac{V}{T} = \text{constant}]$ Ideal Gas Law: $[PV = nRT]$ Where: (P) = pressure (V) = volume (n) = number of moles (R) = universal gas constant (T) = temperature in Kelvin

Oscillations and Waves Describes periodic motions and wave phenomena.

1. Simple Harmonic Motion (SHM) Displacement: $[x(t) = A \sin(\omega t + \phi)]$ Velocity: $[v(t) = A \omega \cos(\omega t + \phi)]$ Acceleration: $[a(t) = -A \omega^2 \sin(\omega t + \phi) = -\omega^2 x(t)]$ Where: (A) = amplitude (ω) = angular frequency (ϕ) = phase constant Period and Frequency: $[T = \frac{2\pi}{\omega}]$ $[f = \frac{1}{T}]$ Restoring Force: $[F = -kx]$ where (k) is the force constant.

2. Wave Speed $[v = \lambda f]$ Where: (λ) = wavelength (f) = frequency

Electromagnetism Covers electric charges, potentials, currents, and magnetic effects.

1. Coulomb's Law $[F = k_e \frac{q_1 q_2}{r^2}]$ Where: (k_e) = Coulomb's constant $(9 \times 10^9 \text{ Nm}^2/\text{C}^2)$ (q_1, q_2) = point charges

2. Ohm's Law $[V = IR]$ Where: (V) = voltage (I) = current (R) = resistance

3. Magnetic Force on a Moving Charge $[F = q v B \sin \theta]$ Where: (q) = charge

QuestionAnswer

What are the main formulas related to kinematics in Class XII physics?

The main kinematic formulas include: $v = u + at$, $s = ut + \frac{1}{2} at^2$, $v^2 = u^2 + 2as$, where u = initial velocity, v = final velocity, a = acceleration, s = displacement, and t = time.

What is the formula for calculating potential energy in physics?

Potential energy (PE) is given by $PE = mgh$, where m is mass, g is acceleration due to gravity, and h is height above the reference point.

How is the formula for Ohm's Law expressed in Class XII physics?

Ohm's Law states that $V = IR$, where V is voltage, I is current, and R is resistance.

What is the formula for calculating work done in physics?

Work done (W) is calculated as $W = Fd \cos \theta$, where F is the applied force, d is the displacement, and θ is the angle between force and displacement.

Can you list the formulas related to Newton's Laws of Motion?

Newton's Second Law is $F = ma$, where F is force, m is mass, and a is acceleration. Also, the law implies that the force acting on an object is equal to the rate of change of its momentum.

Related keywords: physics formulas, class 12 physics, CBSE physics formulas, class 12 formulas list, physics equations class XII, physics derivations class 12, class 12 physics notes, physics concepts class 12, physics problem-solving formulas, class 12 physics syllabus

ICSE class 10 physics numericals

ICSE Class 10 Physics Numericals are an integral part of the curriculum that helps students develop a thorough understanding of fundamental concepts and enhance their problem-solving skills. Numericals are vital for mastering physics as they provide practical application opportunities for theories learned in the classroom. For ICSE Class 10 students, practicing numericals regularly not only boosts confidence but also prepares them efficiently for exams. This article offers a comprehensive guide on ICSE Class 10 physics numericals, including important topics, step-by-step solving techniques, tips, and sample problems to build a solid foundation.

Understanding the Significance of Physics Numericals in ICSE Class 10

Physics is a subject that relies heavily on numerical problems to demonstrate understanding of concepts such as motion, forces, energy, and electricity. Solving numericals:

- Reinforces theoretical concepts
- Develops analytical thinking
- Prepares students for exam pressure
- Improves application skills

In ICSE Class 10, numericals are often based on chapters like Motion, Force and Laws of Motion, Gravity, Work, Power and Energy, Sound, Light, and Electricity. Mastery over these problems is crucial for scoring well in physics papers.

Common Topics Covered in ICSE Class 10 Physics Numericals

To excel in numericals, students must

familiarize themselves with the key chapters and their associated problems:

1. Motion Uniform and non-uniform motion Equations of motion Speed, velocity, and acceleration Distance-time and velocity-time graphs
2. Force and Laws of Motion Newton's laws Friction Momentum and impulse
3. Gravitation Universal law of gravitation Acceleration due to gravity Orbits and satellite motion
4. Work, Power, and Energy Work done Power Kinetic and potential energy Conservation of energy
5. Sound Speed of sound Reflection and echo Noise pollution
6. Light Reflection and refraction Lens formulas Magnification
7. Electricity Ohm's law Resistance, current, voltage Series and parallel circuits Power consumption

Step-by-Step Approach to Solving ICSE Class 10 Physics Numericals Effective problem-solving requires a systematic approach. Here's a step-by-step guide:

1. Read the Problem Carefully Identify what is given and what needs to be found. Note units and relevant quantities.
2. Choose the Appropriate Formula Recall the relevant equations associated with the problem. Write down the formula before substituting values.
3. Convert Units if Necessary Ensure all quantities are in SI units for consistency. Convert units like km/h to m/s, cm to meters, etc.
4. Substitute Values and Calculate Plug in the numerical values carefully. Follow order of operations. Keep track of significant figures.
5. Check the Reasonableness of the Answer Verify the units. Check if the magnitude makes sense. Recalculate if necessary.
6. Write the Final Answer Clearly State the answer with proper units. Highlight the key result.

Sample Numerical Problems and Solutions Below are some illustrative examples that demonstrate typical ICSE Class 10 physics numericals, along with detailed solutions.

Problem 1: Calculating Speed from Distance and Time A car covers a distance of 150 km in 3 hours. Find the speed of the car in km/h and m/s.

Solution: Given: Distance, $(s = 150, \text{ km})$ Time, $(t = 3, \text{ hours})$ To find: Speed in km/h and m/s

Step 1: Calculate speed in km/h
$$\text{Speed} = \frac{\text{Distance}}{\text{Time}} = \frac{150, \text{ km}}{3, \text{ hours}} = 50, \text{ km/h}$$

Step 2: Convert km/h to m/s
$$1, \text{ km/h} = \frac{1000, \text{ m}}{3600, \text{ s}} = \frac{1}{3.6}, \text{ m/s}$$

$$\text{Speed} = 50 \times \frac{1}{3.6} \approx 13.89, \text{ m/s}$$

Answer: Speed = 50 km/h Speed = 13.89 m/s

Problem 2: Calculating Force using Newton's Second Law A mass of 10 kg is accelerated at a rate of 5 m/s². Find the force applied.

Solution: Given: Mass, $(m = 10, \text{ kg})$ Acceleration, $(a = 5, \text{ m/s}^2)$

Formula: $[F = m \times a]$

Calculation: $[F = 10 \times 5 = 50, \text{ N}]$

Answer: Force = 50 Newtons

Problem 3: Calculating Work Done A person lifts a box weighing 20 kg to a height of 5 meters. Calculate the work done in lifting the box. (Take $(g = 9.8, \text{ m/s}^2)$)

Solution: Given: Mass, $(m = 20, \text{ kg})$ Height, $(h = 5, \text{ m})$ Gravitational acceleration, $(g = 9.8, \text{ m/s}^2)$

Formula: $[W = m \times g \times h]$

Calculation: $[W = 20 \times 9.8 \times 5 = 980, \text{ J}]$

Answer: Work done = 980 Joules

Tips for Mastering ICSE Class 10 Physics Numericals To excel in solving numericals, students should adopt effective strategies:

- Practice Regularly:** Solve a variety of problems from textbooks, previous years' papers, and sample papers.
- Understand Concepts:** Grasp the underlying principles before attempting numerical problems.
- Memorize Key Formulas:** Keep a formula sheet handy for quick reference.
- Work on Units:** Always convert quantities into SI units before calculation.
- Check Your Work:** Review calculations, units, and reasonableness of the answer.
- Time Management:** Allocate sufficient time for numericals during exams without rushing.

Additional Resources for ICSE Class 10 Physics Numericals Textbooks and Practice Books: Use NCERT and ICSE-specific physics books with

solved numericals. Sample Papers and Previous Year Questions: Practice past exams for familiarity. Online Tutorials and Video Lectures: Visual explanations can clarify complex problems. Study Groups: Collaborate with peers for doubt clearing and problem-solving practice. Conclusion Mastering ICSE Class 10 physics numericals is crucial for academic success in physics. A disciplined approach, consistent practice, and understanding of concepts are key to solving problems efficiently. By focusing on chapter-wise topics, following systematic methods, and practicing a variety of numerical problems, students can build confidence and excel in their exams. Remember, numericals not only assess your grasp of physics but also prepare you for real-world problem-solving skills essential in scientific pursuits and everyday life. Start practicing today to unlock your potential and achieve excellent results in ICSE Class 10 Physics!

ICSE Class 10 Physics Numericals: An In-depth Analysis for Students and Educators In the realm of ICSE Class 10 Physics, numericals form the backbone of understanding core concepts and applying theoretical knowledge to real-world problems. These numerical questions not only test students' grasp over formulas and principles but also sharpen their analytical thinking and problem-solving skills. Given the importance of these exercises in board examinations, a systematic and comprehensive approach to mastering them can significantly enhance a student's performance. This article aims to delve deep into the nature of ICSE Class 10 Physics numericals, explain key concepts in detail, and provide strategic insights to approach and solve these problems effectively.

The Significance of Numericals in ICSE Class 10 Physics Numerical problems in ICSE Class 10 Physics serve multiple purposes:

- Application of Concepts:** They bridge the gap between theoretical understanding and practical application, ensuring students can practically deploy formulas.
- Assessment of Analytical Skills:** Numericals evaluate how well students can interpret data, manipulate equations, and arrive at logical solutions.
- Preparation for Exams:** Since exams often include numerical questions, practicing these enhances confidence and proficiency.
- Understanding Real-World Physics:** These problems often mimic real-life scenarios, making physics more tangible and relatable.

Given their importance, mastering numericals is critical for excelling in the subject.

Structure and Types of Numericals in ICSE Class 10 Physics Numerical problems in ICSE Class 10 Physics predominantly fall into categories based on the chapter or concept they cover:

- Common Types of Numericals**
 - Mechanical Quantities:** Calculations involving velocity, acceleration, force, work, power, and energy.
 - Optics:** Problems related to refraction, reflection, lens formula, mirror formula, magnification.
 - Sound:** Questions involving speed of sound, echo, and related phenomena.
 - Thermal Physics:** Problems involving heat transfer, specific heat capacity, and thermal expansion.
 - Electricity and Magnetism:** Calculations involving current, voltage, resistance, power, and magnetic effects.
 - Modern Physics:** Numerical based on photoelectric effect, atomic models, and related concepts.

Typical Format of Numerical Questions

Given data (e.g., distances, times, angles, resistances) Required to find (e.g., velocity, refractive index, work done) Use of formulas or principles (e.g., $v = u + at$, $\frac{1}{f} = \frac{1}{v} + \frac{1}{u}$) Step-by-step calculations leading to the final answer

Core Concepts and Formulas for Solving Numericals A thorough understanding of

fundamental formulas is essential. Here's a detailed overview of important equations and principles used in ICSE Class 10 Physics numericals:

Mechanics

Velocity, $(v = \frac{s}{t})$ (s = displacement, t = time)

Acceleration, $(a = \frac{\Delta v}{\Delta t})$

Force, $(F = ma)$ (Newton's Second Law)

Work, $(W = F \times s \times \cos \theta)$

Power, $(P = \frac{W}{t})$

Kinetic Energy, $(KE = \frac{1}{2}mv^2)$

Potential Energy, $(PE = mgh)$

Optics

Mirror formula, $(\frac{1}{u} + \frac{1}{v} = \frac{1}{f})$

Refraction index, $(n = \frac{\sin i}{\sin r})$

Magnification, $(m = \frac{v}{u})$

Sound

Speed of sound, $(v = f \lambda)$ (f = frequency, λ = wavelength)

Thermal Physics

Heat transferred, $(Q = mc \Delta T)$

Thermal expansion, $(\Delta L = \alpha L \Delta T)$

Electricity

Ohm's Law, $(V = IR)$

Power, $(P = VI = I^2 R = \frac{V^2}{R})$

Step-by-Step Approach to Solving Numericals

Mastering numericals requires a strategic approach:

- Carefully Read the Question** Identify what is given and what needs to be found.
- Note units and convert them if necessary.**
- List Known Data and Unknowns** Write down the known quantities. Assign symbols to unknown quantities for clarity.
- Choose the Appropriate Formula** Recall the relevant formula(s) based on the problem. Consider diagrammatic representations if applicable.
- Substitute the Known Values** Plug in the numerical data into the formula.
- Maintain consistent units.**
- Perform Calculation with Precision** Use a calculator for accuracy.
- Keep track of significant figures.**
- Cross-Check Units and Magnitudes** Verify if the answer makes physical sense.
- Check units to avoid errors.**
- Final Write-up** State the answer clearly with proper units. Include necessary steps if required by the examination pattern.

Sample Numerical Problems and Solutions

To illustrate the process, let's analyze some typical numericals encountered in ICSE Class 10 Physics.

Example 1: Calculating Velocity

Question: A car travels a distance of 150 km in 3 hours. Find its average velocity in m/s.

Solution: Given: $(s = 150 \text{ km} = 150,000 \text{ m})$, $(t = 3 \text{ hours} = 3 \times 3600 = 10,800 \text{ s})$

Apply: $(v = \frac{s}{t})$

Calculation: $[v = \frac{150,000}{10,800} \approx 13.89 \text{ m/s}]$

Answer: The average velocity is approximately 13.89 m/s.

Example 2: Refraction and Refractive Index

Question: An incident ray strikes a glass slab at an angle of (30°) and refracts at an angle of (20°) . Find the refractive index of glass.

Solution: Formula: $(n = \frac{\sin i}{\sin r})$

Values: $[n = \frac{\sin 30^\circ}{\sin 20^\circ} = \frac{0.5}{0.3420} \approx 1.46]$

Answer: The refractive index of glass is approximately 1.46.

Example 3: Work Done in Moving an Object

Question: A force of 50 N moves a box through a distance of 10 meters in the direction of the force. Calculate the work done.

Solution: Given: $(F = 50 \text{ N})$, $(s = 10 \text{ m})$, $(\theta = 0^\circ)$ (since movement is in the same direction)

Apply: $(W = F \times s \times \cos \theta)$

$[W = 50 \times 10 \times \cos 0^\circ = 50 \times 10 \times 1 = 500 \text{ J}]$

Answer: The work done is 500 Joules.

Common Challenges and Tips for Mastery

Challenges Faced by Students

- Incorrect Unit Conversion:** Leading to wrong answers.
- Misapplication of Formulas:** Using wrong formulas or mixing concepts.
- Calculation Errors:** Arithmetic mistakes or oversight.
- Time Management:** Spending too long on complex problems.

Tips for Effective Practice

- Master Formulas:** Memorize and understand the derivation and application.
- Practice Regularly:** Solve a variety of problems to build confidence.
- Understand Concepts:** Don't memorize blindly; grasp the physics behind formulas.
- Use Diagrams:** Visual aids help in understanding and solving.
- Review Mistakes:** Analyze errors to avoid repeating them.

Recommended Resources

- ICSE Physics Textbooks:** Focused practice based on prescribed chapters.
- Sample Question Banks:** Past papers and model questions.
- Online Tutorials:** Visual

explanations and step-by-step solutions. Study Groups: Collaborative problem-solving. Conclusion: The Path to Excellence in ICSE Class 10 Physics Numericals Numericals in ICSE Class 10 Physics are not just questions to be answered but opportunities to demonstrate a comprehensive understanding of physical principles. Success hinges on a solid grasp of formulas, logical problem-solving strategies, and consistent practice. By systematically approaching each problem, verifying calculations, and understanding the underlying concepts, students can navigate the numerical section confidently. This mastery not only paves the way for excellent exam scores but also cultivates a deeper appreciation of the fascinating world of physics

QuestionAnswer

What is the formula to calculate the potential energy stored in a stretched wire in ICSE Class 10 Physics numericals?

The potential energy (U) stored in a stretched wire is given by $U = \frac{1}{2} \times k \times x^2$, where k is the force constant (spring constant) and x is the extension or compression.

How do you determine the final velocity of a body sliding down an inclined plane with friction, given initial velocity, incline angle, and friction coefficient?

Use the energy or force analysis: $v^2 = u^2 + 2g s \sin\theta - 2\mu g s \cos\theta$, where u is initial velocity, s is the distance moved, θ is the incline angle, and μ is the coefficient of friction.

In a simple pendulum, how is the time period affected if the length of the pendulum is doubled?

The time period T is proportional to the square root of the length ($T \propto \sqrt{l}$). So, if the length is doubled, T increases by a factor of $\sqrt{2}$.

Calculate the work done in stretching a wire by a small extension when the force-extension graph is a straight line.

The work done $W = \frac{1}{2} \times \text{Force (F)} \times \text{extension (x)}$. If the graph is linear, the area under the graph (a triangle or rectangle) represents work done.

How can you find the speed of a body after it slides down an inclined plane with friction, given the initial potential energy and work done against friction?

Use energy conservation: $v = \sqrt{2(g h - \text{work done against friction})}$, where h is the height corresponding to potential energy, and work done = friction force $\times$ distance.

In an ICSE numericals problem, how do you calculate the elastic potential energy stored in a stretched wire?

Elastic potential energy = $(1/2) \times k \times x^2$, where k is the spring constant and x is the extension or compression.

What is the relation between the period of a simple pendulum and the acceleration due to gravity, and how is it used in numericals?

The period $T = 2\pi \sqrt{l/g}$. In numericals, rearranging gives $g = (4\pi^2 l) / T^2$ to calculate gravity based on measured period and length.

How do you calculate the power developed in an electric circuit numericals involving work done and time?

Power $P = \text{Work done (W)} / \text{Time (t)}$. In numericals, work done can be calculated as $W = \text{Voltage (V)} \times \text{Current (I)} \times \text{time}$ or using energy formulas.

When calculating the speed of sound in a medium using a numerical problem, what key parameters are needed?

You need the distance traveled by sound (d) and the time taken (t). Speed of sound $v = d / t$. The problem may also involve temperature-dependent formulas if applicable.

Related keywords: ICSE physics numericals, class 10 physics problems, ICSE physics solutions, class 10 physics practice, ICSE physics exercises, class 10 physics questions, ICSE physics formulas, physics numerical examples, class 10 physics tips, ICSE physics revision

Important formula of science for class 10

Important Formula of Science for Class 10 Understanding and mastering the important formulas of science is crucial for Class 10 students aiming to excel in their exams. These formulas form the backbone of concepts in Physics, Chemistry, and Biology, helping students solve numerical problems efficiently and grasp fundamental scientific principles. This comprehensive guide provides a structured overview of essential formulas across all three science disciplines, organized for easy reference and effective learning. Physics Formulas for Class 10 Physics is the branch of science

concerned with the nature and properties of matter and energy. The formulas here are vital for solving numerical problems related to motion, force, energy, and more.

1. Motion and Mechanics

Speed (v): $v = \frac{d}{t}$ where d = distance traveled, t = time taken

Average Speed: $\text{Average speed} = \frac{\text{Total distance}}{\text{Total time}}$

Velocity (u, v): Initial velocity = u Final velocity = v

Acceleration (a): $a = \frac{v - u}{t}$

Equation of Motion (for uniform acceleration): $v = u + at$ $s = ut + \frac{1}{2} a t^2$ $v^2 = u^2 + 2as$

Force (F) (Newton's Second Law): $F = ma$ where m = mass, a = acceleration

Momentum (p): $p = mv$

2. Work, Power, and Energy

Work Done (W): $W = F \times d \times \cos \theta$ (when force and displacement are not in the same direction)

Power (P): $P = \frac{W}{t}$

Kinetic Energy (KE): $KE = \frac{1}{2} m v^2$

Potential Energy (PE): $PE = mgh$ where g = acceleration due to gravity, h = height

3. Gravitation

Universal Law of Gravitation: $F = G \frac{m_1 m_2}{r^2}$ where G = gravitational constant, m_1, m_2 = masses, r = distance between centers

Acceleration due to Gravity (g): $g = \frac{GM}{r^2}$ where M = mass of Earth, r = radius of Earth

4. Sound and Waves

Speed of Sound in Air: $v = \lambda \times f$ where λ = wavelength, f = frequency

Chemistry Formulas for Class 10

Chemistry deals with the composition, structure, properties, and reactions of matter. These formulas help in balancing chemical equations and understanding chemical concepts.

1. Mole Concept and Stoichiometry

Mole Definition: 1 mole = 6.022×10^{23} particles (Avogadro's number)

Number of Particles (N): $N = n \times N_A$ where n = number of moles, N_A = Avogadro's number

Molar Mass (M): Expressed in g/mol

Number of Moles (n): $n = \frac{\text{mass (m)}}{\text{molar mass (M)}}$

Mass from Moles: $m = n \times M$

Volume of Gas at STP: 1 mole of gas occupies 22.4 liters

2. Chemical Reactions and Equations

Balancing Chemical Equations: Ensure equal atoms of each element on both sides

Examples: Combustion of methane: $CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O$

Photosynthesis: $6CO_2 + 6H_2O \xrightarrow{\text{light}} C_6H_{12}O_6 + 6O_2$

3. Acids, Bases, and Salts

pH Scale: Acidic solution: $pH < 7$ Neutral solution: $pH = 7$ Basic solution: $pH > 7$

pH Calculation: $pH = -\log [H^+]$

Neutralization Reaction: Acid + Base $\rightarrow$ Salt + Water

4. Chemical Formulas of Common Substances

Water: H_2O Carbon dioxide: CO_2 Sodium chloride: $NaCl$ Sulfuric acid: H_2SO_4 Hydrochloric acid: HCl

Biology Formulas and Concepts for Class 10

Biology focuses on living organisms, their structure, functions, and interactions. While it doesn't involve many formulas, key concepts and ratios are essential for understanding biological processes.

1. Human Body and Circulatory System

Blood Flow Rate: $\text{Flow rate} = \frac{\text{Volume of blood}}{\text{Time}}$

Heart Rate: Typically measured in beats per minute (bpm)

Blood Pressure: Systolic / Diastolic (measured in mm Hg)

2. Photosynthesis and Respiration

Photosynthesis Equation: $6CO_2 + 6H_2O \xrightarrow{\text{light}} C_6H_{12}O_6 + 6O_2$

Respiration: Glucose breakdown: $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O + \text{energy}$

3. Genetics and Inheritance

Punnett Square Ratios: For monohybrid crosses: 1:2:1 (genotypic ratio) 3:1 (phenotypic ratio)

Probability of Inheritance: $P = \frac{\text{Number of favorable outcomes}}{\text{Total outcomes}}$

Additional Tips for Memorizing Important Formulas

Create flashcards with formulas and their derivations. Practice numerical problems regularly. Use diagrams to understand concepts linked with formulas. Group formulas based on topics for better retention.

Conclusion

Mastering the important formulas of science for Class 10 is essential for building a strong foundation in science and performing well in exams. Regular practice,

understanding the underlying concepts, and quick reference to this organized guide can significantly enhance your problem-solving skills and confidence. Remember, formulas are the tools that help you unlock the mysteries of the physical and chemical universe, so keep practicing and stay curious!

Important Formula of Science for Class 10: A Comprehensive Guide

Understanding the important formula of science for class 10 is fundamental for students aiming to excel in their exams and develop a solid foundation in physics, chemistry, and biology. These formulas not only serve as the backbone for solving numerical problems but also deepen conceptual understanding across various chapters. Mastery over these key formulas simplifies complex concepts and enhances problem-solving skills, making them an essential part of your science preparation.

Why Are Important Formulas Essential in Class 10 Science?

Before delving into the formulas themselves, it's vital to recognize why they are so crucial:

- Foundation for Problem Solving:** Many questions in exams are based on applying these formulas.
- Conceptual Clarity:** Formulas often encapsulate fundamental principles, enabling quicker understanding.
- Efficient Revision:** Knowing formulas helps in quick revision before exams.
- Application Across Chapters:** Many formulas are applicable across multiple chapters, reinforcing interconnected concepts.

Categorization of Important Formulas in Class 10 Science

Class 10 Science is divided into three primary subjects: Physics, Chemistry, and Biology. Each has its set of crucial formulas that students must memorize and understand.

Physics: Formulas Related to Motion, Force, Work, and Energy

Physics forms the core of understanding how objects move and interact in the natural world.

Kinematic Equations of Motion

These equations describe the motion of objects under uniform acceleration:

$$v = u + at$$

Final velocity (v) = initial velocity (u) + acceleration (a) × time (t)

$$s = ut + \frac{1}{2}at^2$$

Displacement (s) = initial velocity (u) × time (t) + (1/2) × acceleration (a) × t²

$$v^2 = u^2 + 2as$$

Final velocity squared = initial velocity squared + 2 × acceleration × displacement

Newton's Laws of Motion

$$F = ma$$

Force = mass × acceleration

Gravitational Potential Energy

$$PE = mgh$$

Potential energy = mass × gravitational acceleration × height

Kinetic Energy

$$KE = \frac{1}{2}mv^2$$

Kinetic energy = half × mass × velocity squared

Work, Power, and Energy

Work Done (W)

$$W = F \times d \times \cos\theta$$

Work Done = Force × Displacement × cosθ

Power (P)

$$P = \frac{W}{t}$$

Power (P) = Work Done / Time

Chemistry: Formulas for Atomic Structure, Chemical Reactions, and Periodic Table

Chemistry involves understanding atoms, molecules, and reactions, with several fundamental formulas.

Atomic Number and Atomic Mass

Atomic number (Z): Number of protons in an atom.

Atomic mass (A): Total protons + neutrons.

Mole Concept and Molar Mass

$$n = \frac{m}{M}$$

Number of moles (n) = Given mass / Molar mass

Molarity (M)

$$M = \frac{n}{V}$$

Molarity = Number of moles of solute / Volume of solution in liters

Dilution Formula

$$C_1V_1 = C_2V_2$$

Concentration before dilution (C₁) × Volume before dilution (V₁) = Concentration after dilution (C₂) × Volume after dilution (V₂)

Law of Conservation of Mass

Mass of reactants = Mass of products in a chemical reaction.

Biology: Formulas for Human Body, Genetics, and Cell Biology

Biology focuses on understanding living organisms and their processes through important formulas.

Photosynthesis Equation

$$6CO_2 + 6H_2O \xrightarrow{\text{light}} C_6H_{12}O_6 + 6O_2$$

Carbon dioxide + Water → Glucose + Oxygen

(Note: This is a reaction, not a formula, but understanding mole ratios is essential.)

Human Heart Rate and Blood Flow

Heart Rate (beats per minute)

Blood flow rate =

Cross-sectional area $\times$ Velocity

Genetic Ratios (Mendelian Inheritance) Genotypic ratio = 1:2:1 (for heterozygous cross) Phenotypic ratio = 3:1 (dominant to recessive traits)

How to Memorize and Apply Important Formulas Effectively

Memorization is vital, but understanding the application of formulas ensures better retention and problem-solving capability.

Tips for Mastering Formulas:

- Regular Practice: Solve diverse problems to familiarize yourself with different applications.
- Create Formula Sheets: Maintain concise notes for quick revision.
- Understand Derivations: Know how formulas are derived to grasp their application better.
- Use Mnemonics: Develop memory aids for formulas that are hard to memorize.
- Group Study: Discuss formulas with peers for better retention.

Sample Problems and How to Use Formulas

Physics Example: Problem: A car accelerates from 0 to 20 m/s in 5 seconds. Find its acceleration. Solution: Using $v = u + at$, where: $u = 0$ m/s, $v = 20$ m/s, $t = 5$ s, $a = (v - u)/t = (20 - 0)/5 = 4$ m/s²

Chemistry Example: Problem: How many grams of NaCl are needed to prepare 2 liters of a 0.5 M solution? Solution: Using $n = C \times V$: $n = 0.5$ mol/L $\times$ 2 L = 1 mol. Molar mass of NaCl = 58.5 g/mol. Mass = $n \times$ molar mass = 1 mol $\times$ 58.5 g/mol = 58.5 grams

Biology Example: Problem: If a plant cell has a surface area of 150 mm² and a velocity of 2 mm/min for transport processes, what is the total blood flow rate across the surface? Solution: Flow rate = Cross-sectional area $\times$ velocity = 150 mm² $\times$ 2 mm/min = 300 mm³/min

Final Tips for Success in Class 10 Science

- Consistent Revision: Regularly revisit formulas to keep them fresh.
- Understand Concepts: Don't memorize blindly; understand the principles behind formulas.
- Practice Past Papers: Apply formulas to various questions from previous exams.
- Seek Clarification: Clarify doubts immediately to avoid confusion later.
- Stay Organized: Keep a well-maintained notebook of formulas for quick revision.

Conclusion

Mastering the important formula of science for class 10 is a stepping stone toward achieving academic excellence. These formulas encapsulate core scientific principles and serve as powerful tools to analyze and solve problems efficiently. By understanding their derivation, application, and interconnections, students can build confidence and improve their overall grasp of science. Remember, consistent practice and conceptual clarity are key to transforming these formulas from mere memorization to effective problem-solving skills. Embrace the formulas, and science will become not just a subject but a fascinating journey of discovery.

Question Answer

What is the formula for speed in science for Class 10?

Speed = Distance / Time

What is the formula for acceleration in physics?

Acceleration = Change in velocity / Time taken

How is force calculated according to Newton's second law?

Force = Mass $\times$ Acceleration

What is the formula for work done in physics?

Work Done = Force $\times$ Displacement $\times$ cos?

What is the formula for density in science?

Density = Mass / Volume

What is the formula for potential energy in science?

Potential Energy = mass $\times$ gravity $\times$ height (PE = mgh)

How is power calculated in physics?

Power = Work done / Time taken

What is the formula for pressure in science?

Pressure = Force / Area

Related keywords: science formulas, class 10 science, physics formulas, chemistry formulas, biology formulas, important science equations, class 10 formulas list, science formula chart, essential science formulas, key science concepts

Class 9 physics formulas

Class 9 physics formulas are essential tools that help students understand and solve a wide range of problems in physics. Mastery of these formulas not only enhances conceptual clarity but also boosts problem-solving efficiency, which is crucial for scoring well in exams. In this comprehensive guide, we will explore the key physics formulas covered in Class 9, organized topic-wise for easy understanding and quick revision.

1. Motion and Its Types

Understanding the fundamental concepts of motion is vital in physics. The formulas related to motion help describe how objects move under various circumstances.

1.1. Distance, Displacement, and Speed

Distance (d): The total length of the path traveled by an object.

Displacement (s): The shortest straight-line distance from the initial to the

final position. Speed (v): The rate at which an object covers distance. 1.2. Formulas Speed (v): $v = \frac{d}{t}$ where d = distance traveled, t = time taken. Displacement (s): For uniform motion, similar to distance but in a straight line. Average Speed: $\text{Average speed} = \frac{\text{Total distance}}{\text{Total time}}$ 1.3. Velocity and Uniform Acceleration Velocity (v): Speed with a directional component; vector quantity. Uniform acceleration (a): When the acceleration is constant. 1.4. Formulas Final velocity (v): $v = u + at$ where u = initial velocity, a = acceleration, t = time. Displacement with uniform acceleration: $s = ut + \frac{1}{2}at^2$ Velocity after time t: $v^2 = u^2 + 2as$ 2. Laws of Motion Newton's Laws of Motion form the foundation of classical mechanics and are crucial in understanding how objects behave under applied forces. 2.1. Newton's First Law (Law of Inertia) An object at rest remains at rest, and an object in motion continues in motion with the same speed and in the same direction unless acted upon by an external force. 2.2. Newton's Second Law Mathematical form: $F = ma$ Where F = net force applied on the object (in Newtons, N), m = mass of the object (kg), a = acceleration (m/s²). 2.3. Newton's Third Law For every action, there is an equal and opposite reaction. 3. Gravitation Gravitation explains the force of attraction between objects with mass, such as Earth and the moon. 3.1. Universal Law of Gravitation Formula: $F = G \frac{m_1 m_2}{r^2}$ Where G = universal gravitational constant ($6.674 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$), m₁ and m₂ = masses of two objects, r = distance between their centers. 3.2. Acceleration Due to Gravity (g) Formula: $g = \frac{GM}{R^2}$ Where M = mass of Earth, R = radius of Earth (~6371 km). 3.3. Weight of an Object Formula: $W = mg$ Where m = mass of the object, g = acceleration due to gravity. 4. Work, Power, and Energy These concepts deal with the capacity to do work and the transfer of energy in physical systems. 4.1. Work Formula: $W = F \times d \times \cos \theta$ Where F = force applied, d = displacement, θ = angle between force and displacement. 4.2. Power Formula: $P = \frac{W}{t}$ Power is the rate of doing work, measured in Watts (W). 4.3. Kinetic Energy Formula: $KE = \frac{1}{2}mv^2$ m = mass of the object, v = velocity. 4.4. Potential Energy Formula: $PE = mgh$ h = height above the reference point. 5. Surface Tension and Fluid Mechanics Understanding the behavior of liquids and the forces acting on their surfaces. 5.1. Surface Tension Definition: The force acting on the surface of a liquid, tending to minimize its surface area. Formula for pressure difference across a curved surface: $\Delta P = \frac{2T}{r}$ Where T = surface tension, r = radius of curvature. 5.2. Bernoulli's Theorem For an incompressible, non-viscous fluid in steady flow, the sum of pressure energy, kinetic energy, and potential energy remains constant along a streamline: Formula: $P + \frac{1}{2}\rho v^2 + \rho gh = \text{constant}$ Where P = pressure, ρ = density, v = velocity, h = height. 6. Sound and Its Properties Sound involves wave motion through a medium. 6.1. Speed of Sound In gases: $v = \sqrt{\frac{\gamma R T}{M}}$ Where γ = adiabatic index, R = universal gas constant, T = temperature, M = molar mass. 6.2. Relationship with Frequency and Wavelength Formula: $v = \lambda \times f$ Where v = speed of sound, λ = wavelength, f = frequency. 7. Light and Reflection Optical laws and formulas help understand how light behaves. 7.1. Laws of Reflection The angle of incidence (i) equals the angle of reflection (r): Formula: $i = r$ 7.2. Mirror Formula Formula: $\frac{1}{f} = \frac{1}{v} + \frac{1}{u}$ f = focal length of the mirror, v = image distance, u = object distance. 8. Refraction of Light Understanding how light bends when passing through different media. 8.1. Snell's Law Formula: $n_1 \sin i = n_2 \sin r$ n₁, n₂ = refractive indices of media; i =

angle of incidence; r = angle of refraction. 8.2. Refractive Index

Class 9 Physics Formulas: A Comprehensive Guide for Students

Understanding and memorizing the class 9 physics formulas is essential for students aiming to excel in their physics exams and develop a clear conceptual understanding of the subject. Physics, being a branch of science that deals with the fundamental principles governing the universe, relies heavily on formulas to quantify phenomena such as motion, force, energy, and electricity. This guide aims to provide a detailed breakdown of all the important formulas, their derivations where necessary, and practical tips for mastering them to build a strong foundation in physics.

Why Are Physics Formulas Important?

Before diving into the formulas, it's important to understand their significance:

- Simplify complex problems:** Formulas condense complex concepts into manageable calculations.
- Aid in problem-solving:** Recognizing which formula applies makes solving questions quicker and easier.
- Build conceptual clarity:** Understanding the derivation and application enhances comprehension.
- Prepare for exams:** Memorizing key formulas is crucial for solving numerical questions efficiently.

Class 9 Physics Syllabus Overview

The physics syllabus typically covers:

- Motion & Measurement
- Force & Laws of Motion
- Gravitation
- Work, Energy & Power
- Sound
- Reflection & Refraction
- Human Eye & The Colourful World
- Electricity & Magnetism

Each section involves specific formulas that are fundamental for solving related problems.

Key Physics Formulas in Class 9

Below, we categorize the formulas based on the topics for easy reference.

Motion & Measurement

a. Distance, Speed, and Time

Speed (v): $v = s / t$

Where: v = speed (m/s or km/h) s = distance traveled (meters or kilometers) t = time taken (seconds or hours)

Average Speed (when speed varies): $v_{avg} = (s_1 + s_2 + \dots + s_n) / (t_1 + t_2 + \dots + t_n)$

b. Uniform Velocity and Uniform Acceleration

Velocity (v) after acceleration: $v = u + at$

Where: u = initial velocity a = acceleration t = time

Displacement (s) with uniform acceleration: $s = ut + \frac{1}{2}at^2$

Final velocity squared: $v^2 = u^2 + 2as$

Force & Laws of Motion

a. Force and Its Effects

Newton's Second Law: $F = ma$

Where: F = force (newtons, N) m = mass (kg) a = acceleration (m/s²)

b. Friction

Frictional Force: $F_{friction} = \mu N$

Where: μ = coefficient of friction N = normal force

c. Momentum

Momentum (p): $p = mv$

Impulse (J): $J = F \times t = \Delta p$

Gravitation

Universal Law of Gravitation: $F = G \frac{m_1 m_2}{r^2}$

Where: G = gravitational constant ($6.674 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2$) m_1, m_2 = masses of objects r = distance between centers of masses

Acceleration due to Gravity (g): $g = GM / r^2$

Weight (W): $W = mg$

Work, Energy & Power

Work Done (W): $W = F \times s \times \cos\theta$

Where: θ = angle between force and displacement

Kinetic Energy (KE): $KE = \frac{1}{2}mv^2$

Potential Energy (PE): $PE = mgh$

Power (P): $P = W / t$

Sound

Speed of Sound: $v = \sqrt{B / \rho}$

Where: B = bulk modulus of the medium ρ = density of the medium

Reflection & Refraction

a. Reflection of Light

Laws of Reflection:

- Angle of incidence (i) = angle of reflection (r)
- Incident ray, reflected ray, and normal all lie in the same plane.

b. Refraction of Light

Refraction Law (Snell's Law): $n_1 \sin i = n_2 \sin r$

Where: n_1, n_2 = refractive indices of media i = angle of incidence r = angle of refraction

Refractive Index: $n = c / v$

Where: c = speed of light in vacuum ($\sim 3 \times 10^8 \text{ m/s}$) v = speed of light in the medium

Human Eye & The Colourful World

Magnification (M): $M = v / u$

Where: v = image distance u = object distance

Power of Lens: $P = 1 / f$ (in meters)

Where: P = power in diopters (D) f = focal length

Electricity & Magnetism

a. Electric Current & Resistance

Ohm's

Law: $V = IR$ Where: V = voltage (volts) I = current (amperes) R = resistance (ohms) Power consumed: $P = VI = I^2 R = V^2 / R$ b. Series & Parallel Circuits Total Resistance in Series: $R_{\text{total}} = R_1 + R_2 + R_3 + \dots$ Total Resistance in Parallel: $1 / R_{\text{total}} = 1 / R_1 + 1 / R_2 + 1 / R_3 + \dots$ c. Magnetic Effect of Electric Current Magnetic Field (B) due to a Current-Carrying Conductor (Ampere's Law): $B = (\mu_0 I) / (2r)$ Where: μ_0 = permeability of free space ($4\pi \times 10^{-7} \text{ T}\cdot\text{m/A}$)

Tips for Memorizing and Using Class 9 Physics Formulas

Create a formula sheet: Summarize all formulas in a dedicated notebook for quick revision.

Understand the derivations: Knowing how a formula is derived helps in remembering and applying it correctly.

Practice numerical problems: Regular solving boosts confidence and solidifies concepts.

Use mnemonic devices: For example, $F = ma$ can be remembered as 'Force equals mass times acceleration.'

Relate formulas to real-life examples: Visualizing how a formula applies in daily life aids understanding.

Practical Application and Problem-Solving Strategies

Identify the topic: Determine which section the problem belongs to.

List knowns and unknowns: Write down what information you have and what you need to find.

Select the relevant formula: Match the problem with the correct formula.

Plug in the values carefully: Watch units and conversions.

Solve systematically: Simplify step-by-step and check units for consistency.

Verify your answer: Cross-check with reasonableness and units.

Conclusion

Mastering the class 9 physics formulas is a gateway to understanding the fundamental principles of the physical world. While memorization is important, focusing on the conceptual understanding behind each formula will make problem-solving more intuitive and less daunting. Regular practice, along with clear notes and structured revision, will ensure that these formulas become second nature, setting a solid foundation for higher studies in physics and related sciences. Remember, physics is not just about formulas; it's about understanding the universe around us through logical reasoning and critical thinking. Happy studying!

Question Answer

What are the main formulas for calculating velocity and acceleration in Class 9 Physics?

Velocity (v) = Displacement (s) / Time (t); Acceleration (a) = Change in velocity / Time taken.

How do you calculate the force using Newton's Second Law in Class 9 Physics?

Force (F) = Mass (m) $\times$ Acceleration (a).

What is the formula for work done in physics, and what are its units?

Work done (W) = Force (F) $\times$ Displacement (d) $\times$ $\cos\theta$; Unit: Joule (J).

How is power defined in physics, and what is its formula?

Power (P) = Work done (W) / Time (t).

What is the formula for calculating the density of an object?

Density (?) = Mass (m) / Volume (V).

Related keywords: kinematics formulas, Newton's laws formulas, work and energy formulas, momentum formulas, gravitation formulas, optics formulas, thermodynamics formulas, sound formulas, electricity formulas, magnetism formulas

Up board physics formula for class 12th

up board physics formula for class 12th Understanding the core formulas in physics is essential for students preparing for the UP Board Class 12 examinations. These formulas serve as the foundation for solving numerical problems and understanding conceptual topics effectively. This article provides a comprehensive compilation of all important physics formulas tailored specifically for UP Board Class 12 students. Covering topics from Mechanics to Modern Physics, this guide aims to assist students in their revision and exam preparation.

Mechanics

- Kinematics**
 - Displacement, $s = ut + \frac{1}{2}at^2$
 - Velocity, $v = u + at$
 - Final velocity (relation with initial velocity and displacement), $v^2 = u^2 + 2as$
- Dynamics**
 - Newton's Second Law, $F = ma$
 - Weight, $W = mg$
 - Frictional Force, $F_f = \mu N$ (where N is the normal force)
- Work, Energy, and Power**
 - Work done, $W = F s \cos \theta$
 - Kinetic Energy, $KE = \frac{1}{2}mv^2$
 - Potential Energy, $PE = mgh$
 - Power, $P = \frac{W}{t}$ or $P = F v \cos \theta$
- Center of Mass and Collision**
 - Center of Mass Velocity, $V_{cm} = \frac{\sum m_i v_i}{\sum m_i}$
 - Elastic Collision, $v_{1f} - v_{2f} = -(v_{1i} - v_{2i})$

Oscillations and Waves

- Simple Harmonic Motion (SHM)**
 - Displacement, $x(t) = A \sin(\omega t + \phi)$
 - Angular frequency, $\omega = 2\pi f = \sqrt{\frac{k}{m}}$ (for spring-mass system)
 - Period, $T = \frac{1}{f} = \frac{2\pi}{\omega}$
 - Maximum velocity, $v_{max} = A\omega$
 - Maximum acceleration, $a_{max} = A\omega^2$
- Wave Motion**
 - Wave velocity, $v = \lambda f$
 - Relation between wave speed, tension, and linear mass density (for strings), $v = \sqrt{\frac{T}{\mu}}$

Properties of Matter and Fluid Mechanics

- Elasticity**
 - Hooke's Law, $F = kx$
 - Stress, $\sigma = \frac{F}{A}$
 - Strain, $\text{Strain} = \frac{\Delta l}{l}$
 - Young's Modulus, $Y = \frac{\sigma}{\text{strain}} = \frac{F l}{A \Delta l}$
- Fluid Mechanics**
 - Continuity Equation, $A_1 v_1 = A_2 v_2$
 - Bernoulli's Equation, $P + \frac{1}{2}\rho v^2 + \rho g h = \text{constant}$
 - Viscosity (Poiseuille's Law), $Q = \frac{\pi r^4 \Delta P}{8 \eta l}$

Thermodynamics

- Laws of Thermodynamics**
 - First Law, $\Delta U = Q - W$
 - Work done during expansion or compression, $W = P \Delta V$ (for quasi-static process)
 - Specific Heat, $C =$

$\frac{Q}{m \Delta T}$)2. Ideal Gas Law $PV = nRT$ Number of moles, $(n = \frac{m}{M})$ (where (M) is molar mass) Electrostatics 1. Coulomb's Law Force between two point charges, $(F = k_e \frac{q_1 q_2}{r^2})$ Electric field due to point charge, $(E = \frac{k_e |q|}{r^2})$ 2. Electric Potential and Capacitance Potential due to point charge, $(V = \frac{k_e q}{r})$ Capacitance of a parallel plate capacitor, $(C = \frac{\epsilon_0 A}{d})$ Energy stored in capacitor, $(U = \frac{1}{2} C V^2)$ Magnetism and Electromagnetism 1. Magnetic Forces Force on a moving charge, $(F = q v B \sin \theta)$ Force on current-carrying wire, $(F = B I L \sin \theta)$ 2. Magnetic Field due to a Long Straight Current Carrying Wire Magnetic field, $(B = \frac{\mu_0 I}{2 \pi r})$ 3. Electromagnetic Induction Faraday's Law, $(\mathcal{E} = - \frac{d\Phi_B}{dt})$ Induced emf in a coil, $(\mathcal{E} = - N \frac{d\Phi_B}{dt})$ Modern Physics 1. Photoelectric Effect Einstein's equation, $(KE_{\max} = hf - \phi)$ Threshold frequency, $(f_0 = \frac{\phi}{h})$ 2. Atomic Physics Energy of photon, $(E = hf)$ De Broglie wavelength, $(\lambda = \frac{h}{p})$ Important Tips for UP Board Class 12 Physics Exam Memorize all the formulas and understand their derivations where possible. Practice numerical problems regularly to build confidence. Revise the formulas periodically to retain them effectively. Understand the units associated with each formula for better application. Utilize diagrams to visualize problems and relate them to formulas. This comprehensive

UP Board Physics Formula for Class 12th: A Comprehensive Guide Understanding the core physics formulas is fundamental for success in Class 12 UP Board examinations. These formulas form the backbone of problem-solving and conceptual clarity in topics ranging from mechanics to optics. This detailed guide aims to provide students with an in-depth understanding of the essential physics formulas, their applications, and tips to master them effectively. Introduction to UP Board Class 12 Physics Syllabus Before diving into formulas, it's crucial to understand the scope of the syllabus. UP Board Class 12 Physics encompasses: Physical Units and Measurement Kinematics Laws of Motion Work, Energy, and Power Rotational Motion Gravitation Properties of Bulk Matter Thermodynamics Oscillations Waves and Sound Electrostatics Current Electricity Magnetism and Electromagnetism Optics Modern Physics Each chapter contains fundamental formulas that are essential for solving numerical problems and understanding concepts. Key Physics Formulas for Class 12 UP Board Below is an organized and detailed compilation of formulas categorized by chapters. This comprehensive list will serve as both a quick revision tool and a detailed reference. 1. Physical Quantities and Measurement Speed $(v): (v = \frac{\text{Distance covered (s)}}{\text{Time taken (t)}})$ Average Velocity: $(v_{\text{avg}} = \frac{\Delta x}{\Delta t})$ Instantaneous Velocity: $(v = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t})$ Acceleration $(a): (a = \frac{\Delta v}{\Delta t})$ 2. Kinematics Equations of Motion (for constant acceleration): $(v = u + at)$ $(s = ut + \frac{1}{2}at^2)$ $(v^2 = u^2 + 2as)$ $(s = \frac{(u + v)}{2} \times t)$ Where: (u) = initial velocity (v) = final velocity (a) = acceleration (s) = displacement (t) = time 3. Laws of Motion Newton's Second Law: $(F = ma)$ Newton's Third Law: For every action, there is an equal and opposite reaction. Universal Law of Gravitation: $(F = G \frac{m_1 m_2}{r^2})$ Gravitational Potential Energy: $(U = - G \frac{m_1 m_2}{r})$

Gravitational Force near Earth's Surface: $(F = mg)$

4. Work, Power, and Energy
 Work Done (W): $(W = F \times s \times \cos \theta)$
 Kinetic Energy (KE): $(KE = \frac{1}{2}mv^2)$
 Potential Energy (PE): $(PE = mgh)$
 Power (P): $(P = \frac{W}{t})$
 Work-Energy Theorem: Work done = Change in kinetic energy

5. Rotational Motion
 Angular Displacement (θ) : Radians
 Angular Velocity (ω) : $(\omega = \frac{\Delta \theta}{\Delta t})$
 Angular Acceleration (α) : $(\alpha = \frac{\Delta \omega}{\Delta t})$
 Relation between linear and angular quantities: $(v = r \omega)$ $(a_t = r \alpha)$
 Moment of Inertia (I): For a point mass: $(I = m r^2)$ For a rigid body: depends on shape
 Torque (τ) : $(\tau = I \alpha)$
 Rotational Kinetic Energy: $(KE_{\text{rot}} = \frac{1}{2} I \omega^2)$

6. Gravitation
 Acceleration due to Gravity at height h: $(g' = g \left(1 - \frac{2h}{R}\right))$
 Kepler's Third Law: $(T^2 \propto r^3)$
 Orbital Velocity: $(v = \sqrt{\frac{GM}{r}})$

7. Properties of Bulk Matter
 Stress $(\sigma = \frac{F}{A})$
 Strain: $(\text{Strain} = \frac{\Delta l}{l})$
 Young's Modulus (Y): $(Y = \frac{\text{Stress}}{\text{Strain}} = \frac{\sigma}{\text{Strain}})$
 Bulk Modulus (K): $(K = -V \frac{\Delta P}{\Delta V})$
 Shear Modulus (S): $(S = \frac{\text{Shear Stress}}{\text{Shear Strain}})$

8. Thermodynamics
 First Law of Thermodynamics: $(\Delta U = Q - W)$
 Heat added in isothermal process: $(Q = nRT \ln \frac{V_2}{V_1})$
 Work done in Isothermal Process: $(W = nRT \ln \frac{V_2}{V_1})$
 Coefficient of Linear Expansion: $(\Delta L = \alpha L \Delta T)$
 Coefficient of Volume Expansion: $(\Delta V = \beta V \Delta T)$

9. Oscillations
 Simple Harmonic Motion (SHM): $(x(t) = A \sin(\omega t + \phi))$
 Angular frequency: $(\omega = 2\pi f = \sqrt{\frac{k}{m}})$
 Period of Oscillation: $(T = \frac{2\pi}{\omega})$
 Maximum velocity: $(v_{\text{max}} = A \omega)$
 Maximum acceleration: $(a_{\text{max}} = A \omega^2)$

10. Waves and Sound
 Wave velocity: $(v = f \lambda)$
 Speed of sound in air: $(v = \sqrt{\frac{\gamma R T}{M}})$
 Intensity of sound: $(I = \frac{P}{A})$

11. Electrostatics
 Coulomb's Law: $(F = k \frac{q_1 q_2}{r^2})$
 Electric Field due to point charge: $(E = \frac{k q}{r^2})$
 Electric Potential: $(V = \frac{k q}{r})$
 Capacitance: $(C = \frac{Q}{V})$
 Energy stored in a capacitor: $(U = \frac{1}{2} C V^2)$

12. Current Electricity
 Ohm's Law: $(V = IR)$
 Power in electrical circuit: $(P = VI = I^2 R = \frac{V^2}{R})$
 Resistivity (ρ) relation: $(R = \frac{\rho l}{A})$
 Internal Resistance and emf: $(V = \mathcal{E} - Ir)$

13. Magnetism and Electromagnetism
 Magnetic Force on a moving charge: $(F = q v B \sin \theta)$
 Biot-Savart Law: $(d\vec{B} = \frac{\mu_0}{4\pi} \frac{I d\vec{l} \times \hat{r}}{r^2})$
 Magnetic Field due to a long straight current-carrying wire at distance r: $(B = \frac{\mu_0 I}{2\pi r})$

QuestionAnswer

What is the formula for velocity in the motion equations in class 12 physics?

The formula for velocity in uniformly accelerated motion is $v = u + at$, where u is initial velocity, a is acceleration, and t is time.

How is the work-energy theorem expressed in UP Board Class 12 physics?

The work-energy theorem states that the work done on a body is equal to the change in its kinetic energy, expressed as $W = \Delta K = K_{\text{final}} - K_{\text{initial}}$.

What is the formula for the potential energy stored in a stretched or compressed spring?

The potential energy stored in a spring is given by $U = (1/2) k x^2$, where k is the spring constant and x is the displacement from equilibrium.

How do you calculate the power used in a physics problem involving work done?

Power is calculated as $P = W / t$, where W is work done and t is the time taken.

What is the formula for the elastic potential energy in terms of stress and strain?

Elastic potential energy per unit volume is given by $U = (1/2) \sigma \epsilon$, where σ is stress and ϵ is strain.

How is the period of a simple pendulum derived in UP Board Class 12 physics?

The period T of a simple pendulum is given by $T = 2\pi \sqrt{l / g}$, where l is the length of the pendulum and g is acceleration due to gravity.

What is the formula for the work done by a variable force?

The work done by a variable force $F(x)$ over a displacement from x_1 to x_2 is $W = \int_{x_1}^{x_2} F(x) dx$.

Related keywords: up board physics formulas, class 12 physics formulas, UP board physics notes, UP board physics chapter formulas, class 12 physics equations, UP board science formulas, physics formula list UP board, UP board class 12 physics syllabus, UP board physics important questions, class 12 physics chapter formulas